



<http://www.biodiversitylibrary.org/>

**Proceedings of the American Academy of Arts and Sciences.**

Boston :Metcalf and Co.,1846-1958

<http://www.biodiversitylibrary.org/bibliography/3934>

**v. 3 (1852-1857):** <http://www.biodiversitylibrary.org/item/22230>

Article/Chapter Title: Opuntia

Author(s): Engelmann

Subject(s): Opuntia fusco-atra

Page(s): Page 297

Contributed by: MBLWHOI Library

Sponsored by: MBLWHOI Library

Generated 5 June 2016 12:24 PM

<http://www.biodiversitylibrary.org/pdf4/052208700022230>

This page intentionally left blank.

*O. FUSIFORMIS*, E. & B. l. c.: subprostrata; radicibus elongato-fusiformibus; articulis orbiculatis; pulvillis setas elongatas virescenti-fuscas gerentibus, plerisque vel solum superioribus armatis; aculeis 2-3 gracilibus albidis deflexis seu patentibus; floribus minoribus; stigmatibus 8; bacca ovata; seminibus majusculis subregularibus.

Kansas and Nebraska, in the regions of the Cross-Timbers, from the Canadian to the Big Bend of the Missouri.—Roots elongated tubers  $\frac{1}{2}$ -1 inch in diameter; joints about 3-4 inches long; spines an inch or a little more in length, slenderer and paler than in *O. Rafinesquii*. Flowers 2- $2\frac{1}{2}$  inches in diameter. Seed  $2\frac{3}{4}$  lines wide. This plant has been distributed by me under the name of *Opuntia bulbosa*.

17. *O. FUSCO-ATRA*, E. in P. R. R.: diffusa; articulis orbiculato-obovatis tuberculatis; pulvillis subremotis magnis griseo-tomentosis, inferioribus solum inermibus; setis numerosis robustis longiusculis fuscis; aculeis subsingulis robustis fusco-atris suberectis, altero breviore deflexo saepe adjecto; floris flavi ovario conico pulvillo 12-18 fulvo-villosos et fusco-setosos gerente; stigmatibus 5.

Sterile places in prairies, west of Houston, Texas: fl. May.—The stout brown, or above almost black spines, and the thick bunches of unusually stout brown bristles on the small joints, give this plant a very distinct appearance. Joints  $2\frac{1}{2}$ -3 inches long; pulvilli 6-9 lines apart; bristles 2-3 lines long; spines 1- $1\frac{1}{4}$  inches long, the lower one, when present, about half as long, but hardly less stout. Flower nearly 3 inches in diameter; ovary an inch long, rather slender, its pulvilli covered with long grayish-brown wool, and the upper ones with a few bright-brown bristles.

18. *O. VULGARIS*, Mill.: diffusa, prostrata; radice fibrosa; articulis obovatis seu suborbiculatis crassis late seu pallide viridibus plerumque inermibus; foliis ovatis cuspidatis fere adpressis; pulvillis subremotis parvis subimmersis setas paucas abbreviatas virescenti-stramineas gerentibus; aculeis rarissimis singulis robustis variegatis suberectis; alabastro subgloboso obtuso; ovario clavato pulvillis sub-10 stipato; petalis sub-8; stigmatibus 5; bacca obovata clavata; seminibus regularibus crassis crasse marginatis.

From the southeastern coast of Massachusetts to Georgia and Florida; apparently only in the low countries east and southeast of the Alleghany Mountains, generally not far from the sea-coast: fl. May and June.—Joints 2-4 inches long and 2- $2\frac{1}{2}$  in diameter, rather thick and fleshy. Leaves 2- $2\frac{1}{2}$  lines long, generally appressed, only in