



<http://www.biodiversitylibrary.org/>

Mémoires du Muséum d'histoire naturelle.

Paris :G. Dufour,1815-1832.

<http://www.biodiversitylibrary.org/bibliography/50067>

t. 17 (1828): <http://www.biodiversitylibrary.org/item/108846>

Page(s): Page 118

Contributed by: Smithsonian Institution Libraries

Sponsored by: Biodiversity Heritage Library

Generated 5 August 2012 7:51 PM

<http://www.biodiversitylibrary.org/pdf4/011617500108846>

This page intentionally left blank.

las tuberculorum, aculeis 5-6 rigidis stramineis setaceis, senioribus epidermide secedente exuviatis. ♪ in Mexico. Coulter, n° 38. Caules vix pollicares. Habitus ferè Stapeliæ cespitosæ aculeis omissis.

O. EXUVIATA, ramosa, erecta, teretiuscula, ramis tuberculis compressis irregulariter cristatisve-instructis ferè pentagonis, areolis orbiculatis velutinis ad axillas tuberculorum, aculeis 6-12 stramineis rigidis rectis, senioribus epidermide secedente exuviatis. ♪ in Mexico. Coulter, n° 18. Cactus tunicatus hort. berol. ex ill. Pr. de Salm-Dyck. Truncus pedalis sesqui-poll. crassus.

β. *angustior*, trunco tenuiore, aculeis paucioribus, areolâ angustiore. Coulter, n° 17.

γ. *spinosior*, caule nano, aculeis longioribus crebrioribus spinosissimo.

O. DECIPIENS, erecta, ramosa, viridis, ramis cylindricis basi attenuatis, tuberculis paucis subspiraliter dispositis, areolâ parvâ, aculeis biformibus, uno inferiore maximo patenti-deflexo, cæteris 3-4 minimis setiformibus subradiantibus. ♪ in Mexico. Coulter, n° 20. Folia parva, ovato-oblonga, decidua. Aculeus major, pollicaris, demum epidermide secedente exuviatus, cæteri 1-2 lin. longi. Confer cum Op. imbricata Haw. ex ill. Pr. de Salm-Dyck in litt.

O. KLEINIÆ, erecta, ramosa, cinereo-viridis, ramis erectis cylindricis etuberculatis, fasciculis ordine spirali sinistrorso dispositis, areolâ velutinâ, aculeis biformibus, aliis setosis innumeris ex albedo rufis, uno maximo inferiore patenti-deflexo gracili albido. ♪ in Mexico. Coulter, n° 21. Caulis digiti majoris crassitie, caulem Cacaliæ Kleiniae referens. Folia minima, oblonga, decidua. Aculeus major, pollicaris. Ad priorem sp. accedit.

O. LEPTOCAULIS, erecta, ramosa, ramis cylindricis erectis etuberculatis, fasciculis lineâ spirali sinistrorsâ dispositis, areolâ submentosâ, aculeis biformibus, aliis circiter 3 inferioribus setaceis nigrescentibus patenti-deflexis, cæteris setosis confertis rufescentibus. ♪ in Mexico. Coulter, n° 22. Caulis crassitie digiti minoris. Refert priorem. Specimina duo subemortua video.