



<http://www.biodiversitylibrary.org/>

**Enumeratio diagnostica cactearum hucusque cognitarum /auctore Ludovico
Pfeiffer.**

Berolini : L. Oehmigke, 1837.

<http://www.biodiversitylibrary.org/bibliography/15207>

Item: <http://www.biodiversitylibrary.org/item/51983>

Page(s): Page 170

Contributed by: Missouri Botanical Garden, Peter H. Raven Library
Sponsored by: Missouri Botanical Garden

This page intentionally left blank.

ramorum coccinei, parum expansi, $1\frac{1}{2}$ poll. diam. Ovarium crassum, $1\frac{1}{2}$ poll. longum, tuberculoso-areolatum. Sepala crassa, subulata, purpurea. Petala brevia, rosacea, erecta, subsordide coccinea. Stamina numerosa incurva, antheris albis. Stylus teres, pallide viridis, stigmatibus 8 viridibus.

71. *O. imbricata* DC.

Prodr. III. p. 471. — *Cer. imbricatus* HAW. rev. p. 70.
Pa: Mexico.

O. erecta teretiuscula, non sulcata, superficie porcis sublobulaeformibus varie subimbricatim tessellatis.

Num sequentis varietas, vel eadem?

72. *O. exuviata* DC.

Revue p. 118. — Hort. Dyck. p. 360. — *O. undulata* HORT. — *O. decipiens major* H. VINDOB.

Pa: Mexico.

O. ramosa erecta teretiuscula; ramis tuberculis compressis vel irregulariter cristatis instructis, fere pentagonis; areolis orbiculatis velutinis ad axillas tuberculorum; aculeis 6 — 12 stramineis rigidis rectis, senioribus epidermide secedente exuviatis.

Truncus pedalis, $1\frac{1}{2}$ poll. crassus (DC.) — Species ad *O. imbricatam* proxime accedens, foliolis duplo longioribus, aculeisque numerosioribus ab ea differt (Salm-D. l. c.).

β . *angustior* DC. l. c. — Trunco tenuiore, aculeis paucioribus, areola angustiore.

γ . *spinosior* DC. l. c. — Caule nano, aculeis longioribus, crebrioribus, spinosissimo. — Num *O. tunicata*?

73. *O. tunicata* H. BEROL.

Cereus tunicatus LEHM. (Nov. act. nat. cur. XVI. P. I. p. 319.) — *O. furiosa* WENDL. (catal. h. Herrnh. 1835.)

Pa: Mexico (sec. LEHM. Brasilia merid.).

C. suberectus, articulato-ramosissimus; articulis basi attenuatis tuberculatis; tuberculis apice impressis, lanatis, spiniferis; spinis membrana mobili subdiaphana tunicatis (LEHM.).

O. erecta ramosa; caule ramisque tuberculosis; tuberculis oblongis depressis; aculeis rectis bipollicaribus et tunicatis (WENDL. l. c.).