

Storm-driven maritime dispersal of prickly pear cacti (*Opuntia* species)

Lucas C. Majure, Gary N. Ervin, and Pat Fitzpatrick

Department of Biological Sciences and GeoResources Institute
Mississippi State University, Mississippi State, MS 39762-9652
E-mail: gervin@biology.msstate.edu

METEOROLOGICAL DATA

Hurricane Katrina was an unusually large major hurricane, resulting in a prolonged period of damaging winds and a 17-foot storm surge in the Greenwood Island area on 29 August 2005. The sustained winds peaked at 88 mph (40ms⁻¹) with gusts up to 105 mph (47ms⁻¹). According to an anemometer at Ingalls Shipyard, winds gusts over 100 mph (45ms⁻¹) began at 6AM, associated with a possible outer eyewall. The inner eyewall made landfall between 9AM and 10AM as well as the maximum sustained winds. Time series of the wind data were obtained from the Hurricane Research Division, and are shown in Fig. 4.

To assess the water elevation evolution, the Advanced CIRCulation (ADCIRC) hydrodynamic model⁴ was used to simulate Katrina's storm surge, and a time series was computed (Fig. 4). The model underpredicts the water height, but correctly shows the surge peaking between 10AM and 11AM. In addition, wave heights of 5-10 feet (1.5m to 3.0m) at the islands, and about 2-3 feet (~1m) at the levee system, were superimposed on the surge.

The wind direction (and roughly the water current as well) started out from the east-southeast during the night, then shifted to the southwest in the afternoon. This suggests either the wind or the surge transported broken cactus from the islands to this mainland region. Either mechanism would have provided the 200 N of force necessary to break or dislodge a typical sized cladode of *O. humifusa* (approx. 0.01cm² surface area, with winds of 60 mph [27ms⁻¹] or greater). This would translate to the equivalent of a 20kg mass applied to the cladode, which is more than sufficient for disarticulation of some of the most fibrous *Opuntia* species (e.g., *O. ficus-indica*)². Future trajectory calculations will be performed to test this hypothesis.

CONCLUSIONS

Maximum wind speeds and the forces they created on the barrier islands were of more than adequate strength to have dislodged *Opuntia* cladodes. In fact, entire plants were observed to have been dislodged from the soil and deposited elsewhere on the islands (Fig. 5).

Wind direction data also suggest a high likelihood that the *Opuntia* colonizing the Greenwood Island peninsula could have originated on Petit Bois Island 16km to the southeast, where *Opuntia* were known to exist prior to Hurricane Katrina (Fig. 4).

These results lend support to the hypothesis that storms and maritime transport of *Opuntia* could contribute to long-distance dispersal of these plants – as well as long-distance dispersal of invertebrate herbivores, such as *Cactoblastis* that may inhabit cladodes at the time of transport. However, it presently is unknown whether *Cactoblastis* larvae could survive such a journey; such information would be valuable in planned modeling efforts for oceanic trajectory mapping.

LITERATURE CITED

- Benson, L. 1982. The Cacti of the United States and Canada. Stanford University Press, Stanford, CA.
- Bobich, E. G. and P. S. Nobel. 2001. Biomechanics and anatomy of cladode junctions for two *Opuntia* (Cactaceae) species and their hybrid. *American Journal of Botany* 88: 391-400.
- Frego, K.A. and R.J. Stanforth. 1985. Factors determining the distribution of *Opuntia fragilis* in the boreal forest of southeastern Manitoba. *Canadian Journal of Botany* 63:2377-2382.
- Luettich, R., and J. Westerink. 2007. A parallel Advanced CIRCulation model for oceanic coastal and estuarine waters. Technical report available at <http://www.adcirc.org>.
- Pimenta-Barrios, E., and Rafael F. del Castillo. 2002. Reproductive ecology. In: P.S. Nobel, ed. *Cacti: Biology and Uses*. University of California Press, Berkeley, CA.
- Pinkava, D.J. 2003. *Opuntia*. In: *Flora of North America* Editorial Committee, eds. 1993+. *Flora of North America North of Mexico*. 12+ vols. New York and Oxford. Vol. 4, pp.123-148.
- Radford, A.E., H.E. Ahles, and C.R. Bell. 1968. *Manual of the Vascular Flora of the Carolinas*. University of North Carolina Press, Chapel Hill, North Carolina.
- Rebman, J.P. and D.J. Pinkava. 2001. *Opuntia* cacti of North America - An overview. *Florida Entomologist* 84:474-483.
- Reyes-Aguero, J.A., Aguirre R., J.R., and A. Valiente-Banuet. 2006. Reproductive biology of *Opuntia*. *A review*. *Journal of Arid Environments* 64:549-585.

INTRODUCTION

The majority of *Opuntia* species in the eastern U.S. are found in the Atlantic and Gulf Coastal Plains. These species tend to occur in sandy soils that provide for more xeric environments. This presumably lessens interspecific competition, an important factor for the slow-growing cacti. Several species of *Opuntia* (e.g., *O. humifusa* (Raf.) Raf., *O. pusilla* (Haw.) Haw., and *O. stricta* (Haw.) Haw.) are found in coastal habitats from North Carolina to Texas^{1,8}. These species are particularly abundant on the barrier islands along the Gulf Coast in Mississippi, Alabama, and Florida.

Opuntia are widely known to be dispersed vegetatively by disarticulation of cladodes from the parent plant^{1,8,9}. For example, *Opuntia pusilla* has easily disarticulated cladodes that are armed with retrorsely barbed spines that pierce fur, skin, and other surfaces^{7,1} (Fig. 1). Accordingly, they can be dispersed by animals passing through populations of these plants⁵. The vegetative propagules root readily and form new plants in suitable habitats (Fig. 2). Such vegetative propagation can be one of the main means of reproduction for some *Opuntia*^{1,3,8,6,9}.

Because of the ease with which vegetative fragments of *Opuntia* regenerate, it is likely that cladodes dislodged by storms in coastal areas function in long-distance maritime transport and establishment of new populations. The present work aimed to document one potential example of such dispersal associated with a major hurricane along the US Gulf Coast.

PLANT SURVEYS

Initial plant surveys were conducted July 2004 at Greenwood Island in Jackson County, Mississippi. Greenwood Island is a peninsula adjacent to Bayou Cassote, just west of the Grand Bay National Estuarine Research Reserve and Grand Bay National Wildlife Refuge (Figs. 3 & 4).

The periphery of Greenwood Island grades into salt marsh, which is the most common vegetation cover in the surveyed area. A portion of the interior of Greenwood Island presently is covered by spoils from dredging operations and mainland expansion from the commercial occupants of the eastern half of the peninsula. The peninsula is surrounded by an earthen levee approximately 4.3m (14ft) in height.

No species of *Opuntia* were seen growing in the surveyed area during 2004. The dominant vegetation on the higher areas (oyster shell middens) consisted of *Quercus virginiana*, *Solidago sempervirens*, *Baccharis halimifolia*, *Cissus trifoliata*, *Ampelopsis arborea*, and *Iva frutescens*. Species more common to the peripheral salt marsh were *Juncus roemerianus*, *Solidago sempervirens*, *Borrchia frutescens*, and *Ipomoea sagittata*.

On January 14, 2006, 4.5 months after hurricane Katrina, a second survey was conducted of the same area. Most of the vegetation had been compressed or removed by the force of the storm, and much of the previously surveyed area was occupied by bare soil.

However, an abundance of disarticulated *Opuntia* cladodes were observed, which appeared to have been deposited on the site from hurricane Katrina. Both *O. humifusa* and *O. stricta* were observed. Generally, these plants consisted of individual pads, but occasionally joined cladodes or whole plants were noted.



Figure 1. *Opuntia pusilla* clinging to clothing, illustrating ease of vegetative dispersal in this species.

Figure 2. *Opuntia* cladodes frequently are encountered in wrack along shorelines. Left and center are *Opuntia humifusa* observed in wrack on Horn Island, MS. Far right is an *O. stricta* plant that established in wrack deposited on an oyster shell midden in Grand Bay, MS.



Figure 3. Aerial photograph on the right shows location of Greenwood Island vegetation surveys (orange points) at the base of the earthen levee. No *Opuntia* were present at these points in July 2004.

Lower panel shows location of Greenwood Island, relative to known populations of *Opuntia* at the time Hurricane Katrina made landfall in August 2005 (yellow points).

The points east of Greenwood Island are *Opuntia* on oyster shell middens (Grand Bay, MS).





Figure 4. Water elevation (feet, in red) due to Hurricane Katrina's storm surge as simulated by the Advanced CIRCulation (ADCIRC) model at the locations denoted by red triangles on Horn Island, Petit Bois Island, and the Greenwood Island peninsula. Also shown are observed wind speed (mph, blue) & direction. The winds originally were from the east-southeast at midnight on 29 August, then shifted to the south by the afternoon. Tropical-storm force sustained winds began at 10PM 28 August, and peaked at 9AM 29 August. Wind gusts above 100 mph began at 6AM 29 August (not shown). ADCIRC shows the surge peaked between 10AM and 11PM. The observed surge based on high water marks was between 17.0 and 17.4 feet (5.25m; the ADCIRC model has a low bias).



Figure 5. Entire *Opuntia humifusa* plant trapped in a slash pine (*Pinus ellioti*) on Horn Island, MS (November 26, 2006). Note in the inset photo the new cladode (middle, right of large basal stem) that appeared to have been produced while the plant was suspended from the tree. Hurricane Katrina made landfall more than one year earlier, in August 2005.

ACKNOWLEDGMENTS

This work was funded by the USGS Invasive Species Science Program, the National Biological Information Infrastructure, and the Northern Gulf Institute, WorldWinds, Incorporated (John C. Stennis Space Center, MS) provided meteorological and ADCIRC data used in this analysis.

