

# Effect of Salt Stress, Proline, and Polyamines on Seed Germination of *Opuntia streptacantha*♦

Ana E. Ochoa-Alfaro<sup>1</sup>, Claudia O. Silva-Ortega<sup>1</sup>, Alicia Becerra-Flora<sup>1</sup>,  
Joel Flores-Rivas<sup>2</sup> and Juan F. Jiménez-Bremont<sup>1\*</sup>

<sup>1</sup> División de Biología Molecular,  
Instituto Potosino de Investigación Científica y Tecnológica,  
Camino a la Presa de San José 2055,  
Apartado Postal 3-74 Tangamanga, 78210,  
San Luis Potosí, San Luis Potosí, México.

<sup>2</sup> División de Ciencias Ambientales,  
Instituto Potosino de Investigación Científica y Tecnológica,  
Camino a la Presa de San José 2055,  
Apartado Postal 3-74 Tangamanga, 78210,  
San Luis Potosí, San Luis Potosí, México

\* Author for correspondence; phone: (+52) 444-8-342000; fax: (+52) 444-8-342010,  
email: [jbremont@ipicyt.edu.mx](mailto:jbremont@ipicyt.edu.mx)

## ABSTRACT

In this study we conducted experiments in order to determine whether polyamines (putrescine, spermidine, and spermine) and proline treatments help to break seed dormancy of *Opuntia streptacantha* and also if they facilitate seed germination under salt stress. The results showed that 1mM proline stimulated a better germination percentage as compared to untreated seeds, although the difference was not statistically significant. With respect to polyamines, they did not break seed dormancy. On the other hand, salinity affected seed germination at the higher concentrations (50 and 75 mM NaCl) used, whereas, at 25 mM NaCl, results were similar to the control. The germination percentage in the treatments with proline (1 and 10 mM) and with polyamines (1µM each) diminished significantly when increasing the concentration of NaCl. The exception was spermine treatment, which alleviates seed germination at the higher concentration of NaCl used (75 mM).

**Keywords:** *Opuntia streptacantha*, Polyamines, Proline, Salt stress, Seed germination.

**Abbreviations:** Spd–Spermidine; Spm–Spermine ; PAs–Polyamines; Pro–Proline; Put–Putrescine.

## 1. INTRODUCTION

Seed germination is a process that begins with the uptake of water by the dry seed and it finishes with elongation of the embryo axis (Bewley and Black 1994). This process is affected by adverse factors such as salinity, which is one of the major abiotic stresses that affect plant growth and development, especially in arid and semiarid regions (Neumann 1997, Chinnusamy et al. 2005, Cony et al. 2006). This stress condition could affect the germination of seeds when high osmotic potential is created, which prevents water uptake (osmotic effect) or causes toxic effects of Na<sup>+</sup> and Cl<sup>-</sup> ions (Okçu et al. 2005 and references therein). Plants respond and adapt to stress to survive by complex molecular responses, including the

---

♦ Received 18 October 2007, Accepted 10 July 2008.

accumulation of compatible solutes, the production of protective proteins, and the expression of different sets of genes, which are part of the plant signaling and defense system against high salt stress (Zhu 2001, Sairam and Tyagi 2004). Proline (Pro) is one of the most widely distributed osmolytes in stress conditions not only in plants but also in other organisms (Delauney and Verma 1993; Bartels and Sunkar 2005). Pro has diverse functions, such as stabilization of subcellular structures (proteins and membranes), it functions as a hydroxyl radical scavenger and serves as a source of carbon and nitrogen (Kavi-Kishor et al. 2005). In addition, Pro could act as a component of signal transduction pathways that regulate stress responsive genes (Khedr et al. 2003). Poljakoff-Mayber et al. (1994), suggested a possible role of Pro during seed germination, since they found that dry seeds of *Kosteletzkya virginica* contain a significant amount of betaine but low Pro, but during germination and in the presence of NaCl the betaine content decreased, while the Pro content increased. Furthermore, Pro application in *Zygophyllum simplex* seeds alleviates either the breaking innate seed dormancy or the effect of low salinity on seed germination (Khan and Ungar 1997).

On the other hand, polyamines (PAs) are polycationic compounds of low molecular weight that have been implicated in various physiological processes, such as rhizogenesis, somatic embryogenesis, pollen formation, flowering and initial fruitlet abscission, dormancy, senescence and the response to abiotic and biotic stresses (Bouchereau et al. 1999, Jiménez-Bremont et al. 2006, 2007, Rodríguez-Kessler et al. 2006, 2008). In addition, it has been suggested that PAs may play a role in the accumulation of seed storage proteins and in the seed maturation (Santanen and Simola 1999) and that they help to promote seed germination (Prakash and Prathapasenan 1988, Sińska and Lewandoska 1991, Zeid and Shedeed 2006). PAs biosynthesis increases greatly in response to stresses and its function is presumed to be protective with a role in scavenging free radicals (Ha et al. 1998, Mansour 2000). NaCl salinity causes both ionic imbalance and water stress, creating membrane damage (Tiburcio et al. 1994). PAs prevent uptake of  $\text{Na}^+$ , loss of  $\text{K}^+$ , and leakage of amino acids and electrolytes from plant tissues (Chattopadhyay et al. 2002). It was suggested that PAs contribute to maintaining cellular cation-anion balance and help in the stabilization of the membrane integrity through their low molecular mass and polycationic nature (Smith 1985).

Several reports in Cactaceae have shown that an increase of the salt concentration diminishes the germination percentage (Romero-Schmidt et al. 1992, Nolasco et al. 1996, Vega-Villasante et al. 1996). In the case of *Opuntia* species, there are studies about chemical and mechanical pretreatments used to stimulate seed germination and thus break seed dormancy (Olvera-Carrillo et al. 2003, Mandujano et al. 2005, 2007, Altare et al. 2006). However, it is not known if PAs and Pro exogenous application breaks seed dormancy or if they protect seed germination under salt stress conditions. For this reason, in this present study we examined the effect of PAs such as putrescine (Put), spermidine (Spd) and spermine (Spm), and Pro in breaking seed dormancy and also if they facilitate seed germination of *Opuntia streptacantha* under salt stress.

## 2. MATERIALS AND METHODS

### 2.1 Material of seeds and germination conditions

*Opuntia streptacantha* seeds were collected of “Cardona” variety cactus pears, of the region of Villa de Zaragoza of the state of San Luis Potosí, México (Silva-Ortega et al. 2008). Seeds collection was January 2005, and they were stored at room temperature during twenty-four months. Seeds were sterilized with 30% (v/V) commercial sodium hypochloride solution for 30 min and with 70% (v/V) ethanol for 4 min, and rinsed three times in sterile distilled water between each treatment.

## 2.2 Treatments of seeds

Seeds were subjected to the following treatments: (a) PAs (1  $\mu$ M Put, Spd or Spm), (b) Pro (1 or 10 mM), (c) and for salt treatment (25, 50, or 75 mM NaCl). We also used mixtures of the previous treatments: (d) Mix of PAs (Put, Spd, and Spm) with each Pro concentration, (e) PAs (Put, Spd, and Spm) with each salt solutions and (f) each Pro concentration with each salt solutions.

For each treatment five replicates, with twenty seeds per replicate, were used. The seeds were placed in Petri dishes containing a perlite support. For the control we added 20 ml of sterile distilled water and for the treatments we added 20 ml of each of the solutions previously described. These experiments were carried out in a growth chamber at 25°C with a photoperiod of 12 h light/12 h dark cycle. The number of seeds germinated was recorded every two days during 2 months, considering that a seed was germinated when it showed the radicle emerged.

## 2.3 Statistical analysis

To explore potential differences in seed germination percentage, a one-way ANOVA was used, with treatment as the main factor. Before analysis, the data were transformed using square-root arcsine, as it is recommended for the percentage data (Sokal and Rohlf 1995). Differences among treatments were explored through orthogonal contrasts.

# 3. RESULTS AND DISCUSSION

## 3.1 Breaking seed dormancy

It has been found that many Cactaceae have prolonged seed dormancy, with low or null germination (Rojas-Aréchiga and Vázquez-Yanes 2000, Flores et al. 2005, 2006, 2008). Specifically, one genus of this family considered as having seed dormancy is *Opuntia*. The first species found to have dormant seeds was *Opuntia aurantiaca* (Archibald 1939) and, up to the present, seed dormancy has been found in 28 *Opuntia* species (see Table 1).

Here we show that in the untreated seeds of *O. streptacantha* the germination percentage was low (14%), meaning that this cactus species presents seed dormancy (Figure 1A). This result is similar to that reported for seeds of other species of the *Opuntia* genus (Table 1). Several dormancy-breaking treatments, including scarification with sulfuric acid, mechanical scarification, after-ripening, cold stratification, leaching, passage through the digestive tract of animals, soaking in hot water, and some combinations of these treatments have been suggested to enhance germination of *Opuntia* seeds. However, these treatments do not always promote germination and, in some cases, treated seeds have germinated to lower percentages than those in the control (Orozco-Segovia et al. 2007). Gibberellic acid (GA<sub>3</sub>) alone seems to be only moderately effective in promoting seed germination of *Opuntia* spp. (Wang et al. 1996). However, it induces a high percentage of germination in combination with other treatments such as rinsing/washing, chemical or mechanical scarification (Orozco-Segovia et al. 2007).

## 3.2 Effect of the application of proline and of polyamines on the germination.

Due to the low germination percentage obtained for *O. streptacantha*, in the present study we evaluated the effect of exogenous application of Pro and PAs on breaking seed dormancy. Our results showed that at 1 mM Pro treatment the germination percentage was higher than the control; however, this difference was not significant ( $F = 1.500$ ,  $P > 0.5$ ). Also, at 10 mM Pro, we observed that there was not a significant difference in the germination percentage with respect to the control ( $F = 0.001$ ,  $P > 0.5$ ; Figure 1A). It has

been reported that Pro treatments (1 and 0.1 mM) alleviated the innate dormancy of *Zygophyllum simplex* seeds and germination reached 40-60% compared to 12% in the control (Khan and Ungar 1997).

The PA treatments used in this experiment did not stimulate the breaking seed dormancy, because there were not significant differences in germination percentages among PAs and the control (Figure 1B). It was observed that with Put treatment the germination percentage was slightly higher than the control, but this difference was not significant ( $F = 0.004$ ,  $P > 0.5$ ). It has been reported that Put promotes the germination of various species, for example *Oryza sativa* (Prakash and Pratapasenan 1988), *Hordeum vulgare* (Locke et al. 2000), *Citrullus lanatus* (Korkmaz et al. 2004), *Cucumis melo* (Korkmaz et al. 2005) and *Medicago sativa* (Zeid and Shedeed 2006). However, other authors have found that Put inhibits germination. Amoroso-Botelho and Juliano-Gualtieri (2001), reported that the treatment with 0.5 mM Put did not affect the seed germination of *Peltophorum dubium*, while a negative effect was observed at 1.0 mM Put. Moreover, at 0.1  $\mu$ M and 0.1 mM Put, the germination of the cacti *Turbinicarpus lophophoroides* and *Turbinicarpus pseudopectinatus* was inhibited (Flores et al. 2008). In other plant species, an effect of PAs in the germination has been reported, for example in apple (*Malus domestica*) the exogenous application of Put and Spd (0.1 and 1 mM) stimulated the germination, whereas Spm inhibited it (Sińska and Lewandowska 1991). In *Peltophorum dubium* Spd treatments (0.5 mM and 1.0 mM) did not affect seed germination (Amoroso-Botelho and Juliano-Gualtieri 2001). Thus, we suggest that the effect of PAs in the seed germination depends on the plant species.

We carried out an experiment with the mixture of Pas (1  $\mu$ M Put, Spd, and Spm) with Pro concentrations (1 and 10 mM). Using the higher concentration of Pro (10 mM) with a mix of PAs (1  $\mu$ M of each one) a negative effect on seed germination was observed, although it was not statistically significant ( $F = 3.025$ ,  $P > 0.05$ ). The lower concentration of Pro (1 mM) and a mix of PAs (1  $\mu$ M of each one) showed a similar behavior than the control ( $F = 0.446$ ,  $P > 0.05$ ; Figure 1A).

### 3.3 Effect of salt stress on the germination.

In a general way, we found that the germination of seeds of *O. streptacantha* was significantly affected by salt stress. The percentage of germination of *O. streptacantha* seeds treated with 25 mM NaCl was similar to the control ( $F = 0.045$ ,  $P > 0.05$ ), while at 50 mM NaCl the germination percentage was lower than the control ( $F = 17.633$ ,  $P < 0.0001$ ) and at 75 mM there was a complete inhibition (Figure 2). The results of this study showed that *O. streptacantha* seed germination is affected by salt stress like in other cacti species. For example, in the “cactus barril” (*Ferocactus peninsulae*), the “cardón gigante” (*Pachycereus pringlei*) and the “cardón barbón” (*Pachycereus pecten-aboriginum*) the seeds were able to germinate at 25 and 50 mM NaCl, while at 100 mM the percentage of germination diminished and at 200 and 300 mM there was a complete inhibition (Romero-Schmidt et al. 1992, Nolasco et al. 1996, Vega-Villasante et al. 1996).

### 3.4 Effect of the application of proline during the germination of seeds under salt stress.

In general, the germination percentage in the treatments with Pro (1 and 10 mM) significantly diminished when increasing the NaCl concentration, leading to an inhibition of germination at the higher concentration (75 mM) (Figure 3). Our results are similar to those reported by Poljakoff-Mayber *et al.* (1994) in *Kosteletzkya virginica*, in which the exogenous application of Pro (10 mM) did not improve germination under salt stress conditions (100 mM NaCl). Khan and Ungar (1997) reported that Pro application in *Zygophyllum simplex* seeds alleviates the effect of low salinity (25 mM NaCl), and induced seed germination, being more effective at the lower Pro concentration (0.1mM) than at the higher concentration (1mM). However, at higher salinity treatments (75 and 125 mM NaCl) Pro was ineffective in alleviating the dormancy of seeds. On the other hand, although in our study the exogenous application of Pro did not improve seed germination under stress, it is important to mention that we have shown a

significant Pro accumulation in cladodes of the same species (*O. streptacantha*) under salt stress (Silva-Ortega et al. 2008), suggesting that the effect of Pro depends on the developmental stage of the plant.

### 3.5 Effect of the application of polyamines during the germination of seeds under salt stress.

The application of exogenous PAs (Put, Spd, or Spm) did not produce a significant difference in the percentage of germination respecting control seeds as described above (Figure 4). In the case of simultaneous application of each PAs and salt stress treatments, we observed that the germination percentage significantly diminished as the salt concentration increased together with the addition of Put and Spd. However, Spm treatment alleviated seed germination (4%) at the higher concentration of NaCl used (75mM); it showed a recovery of 28.57%, as compared to the percentage of germination of control plants (14%). At this concentration of NaCl, seed germination was completely inhibited by salt stress, whereas at these same conditions the application of Put and Spd, or Pro did not stimulate germination.

Our results are in agreement, specifically Spm, with the idea that polyamines may be involved in salt stress protection. In a previous study, we have observed an accumulation of Spm in a bean tolerant cultivar at higher concentrations of NaCl used (150 and 400 mM), suggesting that Spm might be part of the mechanism conferring salt tolerance (unpublished data). Zapata et al. (2004) reported a significant increase in Spm and Spd levels during germination of different plant species such as lettuce (*Lactuca sativa* L.), pepper (*Capsicum annuum*) and broccoli (*Brassica oleraceae* L. var. *Italica* Plenck.) under conditions of salt stress (150 mM NaCl). A protective role for Spm in abiotic stress such as high salt or drought stress has been shown in the *Arabidopsis* spermidine synthase (*acl5/spms*) double mutant that does not produce Spm. This *Arabidopsis* Spm-deficient mutant is hypersensitive to high concentrations of NaCl and also to drought stress; and this phenotype was rescued by exogenously applied Spm (Yamaguchi et al. 2006, 2007). This mutant caused a defect of  $Ca^{+}$  homeostasis, evidencing a protective role for Spm in the context of responses to ionic stress (Yamaguchi et al. 2006). On the other hand, Zhu et al. (2006) reported that the exogenous application of PAs, specifically Spd inhibits  $Na^{+}$  transport from roots to shoots in barley seedlings under conditions of high salinity, suggesting that PAs could be a mechanism for alleviating salt stress. In a recent report, Zhao et al. (2008) showed that exogenous application of PAs, specifically Spd blocked the inward  $Na^{+}$  and  $K^{+}$  currents (especially  $Na^{+}$  currents) in barley root epidermal and cortical cells. In root xylem parenchyma, the inward  $K^{+}$  current was also blocked by extracellular Spd, while the outward  $K^{+}$  current was enhanced. Thus, by repressing  $Na^{+}$  influx into roots and by preventing  $K^{+}$  loss from shoots, PAs improved  $K^{+}/Na^{+}$  homeostasis in barley seedlings to cope with high saline conditions.

## 4. CONCLUSIONS

1. *Opuntia streptacantha* presents seed dormancy, which could not be broken down by the different treatments used (Pro, Pas, and Pro plus PAs).
2. The treatments with higher salinity (50 and 75 mM NaCl) affected dramatically the seed germination, whereas the treatment of 25 mM NaCl allowed a similar germination percentage as the control.
3. Germination under salt stress conditions could not be alleviated by the application of different treatments (Pro and PAs), excepting by Spm which promoted 4% of seed germination at 75 mM NaCl.

## 5. REFERENCES

- Altare, M., S. Trione, J.C. Guevara, and M. Cony. 2006. Stimulation and promotion of germination in *Opuntia ficus-indica* seeds. *Journal of the Professional Association for Cactus Development* 8:91-100.
- Amoroso-Botelho, B., and S.C. Juliano-Gualtieri. 2001. Growth regulators and water stress in canafistula seed germination. *Scientia Agricola* 58:43-49.
- Archibald, E.E.A. 1939. The development of the ovule and seeds of jointed cactus (*Opuntia aurantiaca*). *South African Journal of Science* 36:195-211.
- Bartels, D., and R. Sunkar. 2005. Drought and salt tolerance in plants. *Critical Reviews in Plant Sciences* 24:23-58.
- Baskin, J.M., and C.C. Baskin. 1977. Seed and seedling ecology of *Opuntia compressa* in Tennessee cedar glades. *Journal of the Tennessee Academy of Science* 52:118-122.
- Bewley, J.D., and M. Black. 1994. *Seeds: Physiology of Development and Germination*. 2nd edition. Springer, 460 pp.
- Bouchereau, A., A. Aziz, F. Larher, and J. Martin-Tanguy. 1999. Polyamines and environmental challenges: recent development. *Plant Science* 140:103-125.
- Chattopadhyay, M.K., B.S. Tiwari, G. Chattopadhyay, A. Bose, D.N. Sengupta, and B. Ghosh. 2002. Protective role of exogenous polyamines on salinity-stressed rice (*Oryza sativa*) plants. *Physiologia Plantarum*. 116:192-199.
- Chinnusamy, V., A. Jagendorf, and J.K. Zhu. 2005. Understanding and improving salt tolerance in plants. *Crop Science Society of America* 45:437-448.
- Cony, M.A., S.O. Trione, and J.C. Guevara. 2006. Macrophysiological responses of two forage *Opuntia* species to salt stress. *Journal of the Professional Association for Cactus Development* 8:52-62.
- Cook, C.W. 1942. Insects and weather as they influence growth of cactus on the Central Great Plains. *Ecology* 23:209-214.
- D'Aubeterre, R., Z. Pinero, E. García, and M.A. Figarella. 2006. Efecto de diferentes métodos de escarificación sobre la germinación de cinco especies de cactáceas (*Opuntia ficus-indica*, *Pilosocereus moritzianus*, *Stenocereus griseus*, *Cereus deficiens* y *Cereus hexagonus*) del Estado Lara. Simposio-Taller: Experiencias en Agroforestería ejecutadas o en proceso por el Instituto Nacional de Investigaciones Agrícolas (INIA) 1: 13-17
- Delauney, A.J., and D.P.S. Verma. 1993. Proline biosynthesis and osmoregulation in plants. *Plant Journal* 4:215-223.
- Flores, J., A. Arredondo, and E. Jurado. 2005. Comparative seed germination in species of *Turbinicarpus*: An endangered cacti genus. *Natural Areas Journal* 25:183-187.
- Flores, J., E. Jurado, and A. Arredondo. 2006. Effect of light on germination of seeds of Cactaceae from the Chihuahuan Desert, México. *Seed Science Research* 16:149-155.

- Flores, J., E. Jurado, and J.F. Jiménez-Bremont. 2008. Breaking seed dormancy in specially protected *Turbincarpus lophophoroides* and *Turbincarpus pseudopectinatus* (Cactaceae). *Plant Species Biology* 23:44-47.
- Godínez-Alvarez, H., and A. Valiente-Banuet. 1998. Germination and early seedling growth of Tehuacan Valley cacti species: the role of soils and seed ingestion by dispersers on seedling growth. *Journal of Arid Environments* 39:21-31.
- Ha, H.C., N.S. Sirisoma, P. Kuppasamy, J.L. Zweier, P.M. Woster, and R.A. Casero. 1998. The natural polyamine spermine functions directly as a free radical scavenger. *Proceedings of the National Academy of Sciences of the United States of America* 95:11140-11145.
- Jiménez-Bremont, J.F., A. Becerra-Flora, E. Hernández-Lucero, M. Rodríguez-Kessler, J.A. Acosta-Gallegos, and J.G. Ramírez-Pimentel. 2006. Proline accumulation in two bean cultivars under salt stress and the effect of polyamines and ornithine. *Biologia Plantarum* 50:763-766.
- Jiménez-Bremont, J.F., O.A. Ruiz, and M. Rodríguez-Kessler. 2007. Modulation of spermine levels in maize seedlings subjected to long-term salt stress. *Plant Physiology and Biochemistry*. 45:812-821.
- Kavi-Kishor, P.B., S. Sangam, R.N. Amrutha, P. Sri Laxmi, K.R. Naidu, K.R.S.S. Rao, S. Rao, K.J. Reddy, P. Theriappan, and N. Sreenivasulu. 2005. Regulation of proline biosynthesis, degradation, uptake and transport in higher plants: Its implications in plant growth and abiotic stress tolerance. *Current Science* 88:424-438.
- Khan, M.J., and I.A. Ungar. 1997. Alleviation of seed dormancy in the desert forb *Zygophyllum simplex* L. from Pakistan. *Annals of Botany* 80:395-400.
- Khedr, A.H.A, M.A. Abbas, A.A.A. Wahid, W.P. Quick, and G.M. Abogadallah. 2003. Proline induces the expression of salt-stress-responsive proteins and may improve the adaptation of *Pancreaticum maritimum* L. to salt-stress. *Journal of Experimental Botany* 54:2553-2562.
- Korkmaz, A., I. Tiryaki, M.N. Nas, and N. Ozbay. 2004. Inclusion of plant growth regulators into priming solution improves low-temperature germination and emergence of watermelon seeds. *Canadian Journal of Plant Science* 84:1161-1165.
- Korkmaz, A., N. Ozbay, I. Tiryaki, and M.N. Nas. 2005. Combining priming and plant growth regulators improves muskmelon germination and emergence at low temperatures. *European Journal of Horticultural Science* 70:29-34.
- Locke, J.M., J.H. Bryce, and P.C. Morris. 2000. Contrasting effects of ethylene perception and biosynthesis inhibitors on germination and seedling growth of barley (*Hordeum vulgare* L.). *Journal of Experimental Botany* 51:1843-1849.
- Lotter, W.D., L. Thatcher, L. Rossow, and C.F. Reinhardt. 1999. The influence of baboon predation and time in water on germination and early establishment of *Opuntia stricta* (Australian Pest Pear) in the Kruger National Park. *Koedoe* 42:43-50.
- Mandujano, M.C., J. Golubov, and C. Montaña. 1997. Dormancy and endozoochorous dispersal of *Opuntia rastrera* seeds in the southern Chihuahuan Desert. *Journal of Arid Environments* 36:259-266

- Mandujano, M.C., C. Montaña, and M. Rojas-Aréchiga. 2005. Breaking seed dormancy in *Opuntia rastrera* from the Chihuahuan desert. *Journal of Arid Environments* 62:15-21.
- Mandujano, M.C., J. Golubov, and M. Rojas-Aréchiga. 2007. Efecto del ácido giberélico en la germinación de tres especies del género *Opuntia* (Cactaceae) del Desierto Chihuahuense. *Cactáceas y Suculentas Mexicanas* 52:46-52.
- Mansour, M.M.F. 2000. Nitrogen containing compounds and adaptation of plants to salinity stress. *Biologia Plantarum*. 43:491-500.
- Neumann P. 1997. Salinity resistance and plant growth revisited. *Plant, Cell and Environment* 20:1193-1198.
- Nolasco, H., F. Vega-Villasante, H.L. Romero-Schmidt, and A. Díaz-Rondero. 1996. The effects of salinity, acidity, light and temperature on the germination of seeds of cardón (*Pachycereus pringlei* (S.Wats) Britton & Rose, Cactaceae). *Journal of Arid Environments* 33:87-94.
- Okçu, G., M. D. Kaya, and M. Atak. 2005. Effects of salt and drought stresses on germination and seedling growth of pea (*Pisum sativum* L.). *Turkish Journal of Agriculture and Forestry* 29:237-242.
- Olivares, O.J., H.A. Chimal, L. Scheinvar, S.D. Montiel, B.P. Zavaleta, and E.G. Ramos 1999. Germinación de semillas de xoconostle duraznillo (*Opuntia leucotricha*), de la región de Fresnillo, Zacatecas. In: Aguirre, R.J.R., and J.A. Reyes-Agüero (Eds.), Memoria del VIII Congreso Nacional y III Internacional sobre el Conocimiento y Aprovechamiento del Nopal, pp. 32-33, UASLP, San Luis Potosí, México.
- Olvera-Carrillo, Y., J. Márquez-Guzmán, V.L. Barradas, M.E. Sánchez-Coronado, and A. Orozco-Segovia. 2003. Germination of the hard seed coated *Opuntia tomentosa* SD, a cacti from the México valley. *Journal of Arid Environments* 55:29-42.
- Orozco-Segovia, A., J. Márquez-Guzmán, M.E. Sánchez-Coronado, A. Gamboa De Buen, J.M. Baskin, and C.C. Baskin 2007. Seed anatomy and water uptake in relation to seed dormancy in *Opuntia tomentosa* (Cactaceae, Opuntioideae). *Annals of Botany* 99:581-592.
- Pendley, G.K. 2001. Seed germination experiments in *Opuntia* (Cactaceae) of the Northern Chihuahuan Desert. *Haseltonia* 8:42-50.
- Pilcher, B.L. 1970. Germination of seeds of four species of *Opuntia*. *Cactus and Succulent Journal* 42:281-282.
- Potter, R.L., J.L. Petersen, and D.N. Ueckert. 1984. Germination responses of *Opuntia* spp. to temperature, scarification and other seed treatments. *Weed Science* 32:106-110.
- Prakash, L., and G. Prathapasenan. 1988. Putrescine reduces NaCl-induced inhibition of germination and early seedling growth of rice (*Oryza sativa* L.). *Australian Journal of Plant Physiology* 15:761-767.
- Poljakoff-Mayber, A., G.F. Somers, E. Werker, and J.L. Gallagher. 1994. Seeds of *Kosteletzkya virginica* (Malvaceae): Their structure, germination, and salt tolerance. II germination and salt tolerance. *American Journal of Botany* 81:54-59.



- Reinhardt, C.F., L. Rossouw, L. Thatcher, and W.D. Lotter. 1999. Seed germination of *Opuntia stricta*: implications for management strategies in the Kruger National Park. *South African Journal of Botany* 65:295-298.
- Rodríguez-Kessler, M., A. G. Alpuche-Solís, O.A. Ruiz, and J.F. Jiménez-Bremont. 2006. Effect of salt stress on the regulation of maize (*Zea mays* L.) genes involved in polyamine biosynthesis. *Plant Growth Regulation* 48:175-185.
- Rodríguez-Kessler M., O.A. Ruiz, S. Maiale, J. Ruiz-Herrera, and J.F. Jiménez-Bremont. 2008. Polyamine metabolism in maize tumors induced by *Ustilago maydis*. *Plant Physiology and Biochemistry*. In press, doi:10.1016/j.plaphy.2008.05.012.
- Rojas-Aréchiga, M., and C. Vázquez-Yanes. 2000. Cactus seed germination: A review. *Journal of Arid Environments* 44:85-104.
- Romero-Schmidt, H.L., F. Vega-Villasante, H. Nolasco, and C. Montaña. 1992. The effect of darkness, freezing, acidity and salinity on seed germination of *Ferocactus peninsulae* (Cactaceae). *Journal of Arid Environments* 23:389-395.
- Sairam, R.K., and A. Tyagi. 2004. Physiology and molecular biology of salinity stress tolerance in plants. *Current Science* 86:407-418.
- Sánchez-Venegas, G. 1997. Germinación, viabilidad y características distintivas de la semilla de *Opuntia joconostle* Weber, forma Cuaresmero. *Cactáceas y Suculentas Mexicanas* 42:16-21.
- Santanen, A., and L.K. Simola. 1999. Metabolism of L[U-<sup>14</sup>C]-arginine and L[U-<sup>14</sup>C]-ornithine in maturing and vernalised embryos and megagametophytes of *Picea abies*. *Physiologia Plantarum* 107:433-440.
- Sińska, I., and U. Lewandowska. 1991. Polyamines and ethylene in the removal of embryonal dormancy in apple seeds. *Physiologia Plantarum* 81:59-64.
- Silva-Ortega, C.O., A.E Ochoa-Alfaro, J.A. Reyes-Agüero, G.A. Aguado-Santacruz, and J.F. Jiménez-Bremont. 2008. Salt stress increases the expression of P5CS gene and induces proline accumulation in cactus pear. *Plant Physiology and Biochemistry* 46:82-92.
- Silverstein, R.P. 2005. Germination of native and exotic plant seeds dispersed by coyotes (*Canis latrans*) in Southern California. *Southwestern Naturalist* 50:472-478.
- Smith, T.A. 1985. Polyamines. *Annual Review Plant Physiology* 36:117-143.
- Smreciu, E.A., R.S. Currah, and E. Toop. 1988. Viability and germination of herbaceous perennial species native to Southern Alberta grasslands. *Canadian Field Naturalist* 102:31-38.
- Sokal, R. R., and J. H. Rohlf. 1995. *Biometry: the principles and practice of statistics in biological research*, 3rd edition. W.H. Freeman, New York. 887 pp.
- Tiburcio, A.F., R.T. Besford, T. Capell, A. Borrell, P.S. Testillano, and M.C. Risueno. 1994. Mechanism of polyamine action during senescence responses induced by osmotic stress. *Journal of Experimental Botany* 45:1789-1800.

Timmons, F.L. 1942. The dissemination of prickly pear seed by jack rabbits. *Journal of the American Society of Agronomy* 34:513-520.

Vega-Villasante, F., H. Nolasco, C. Montaña, H. Romero-Schmidt, and E. Vega-Villasante. 1996. Efecto de la temperatura, acidez, iluminación, salinidad, irradiación solar y humedad sobre la germinación de semillas de *Pachycereus pecten-aboriginum* “cardón barbón” (Cactaceae). *Cactáceas y Suculentas Mexicanas* 41:51-60.

Wang, X., P. Felker, A. Paterson, Y. Mizrahi, A. Nerd, and C. Mondragón-Jacobo. 1996. Cross-hybridization and seed germination in *Opuntia* species. *Journal of the Professional Association for Cactus Development* 1:49-60.

Wiggins, I.L., and D.W. Focht. 1967. Seeds and seedlings of *Opuntia echios* J. T. Howell var. *gigantea* Dawson. *Cactus and Succulent Journal* 39:26-30.

Yamaguchi, K., Y. Takahashi, T. Berberich, A. Imai, A. Miyazaki, T. Takahashi, A.J. Michael, and T. Kusano. 2006. The polyamine spermine protects against high salt stress in *Arabidopsis thaliana*. *FEBS Letters* 580:6783-6788.

Yamaguchi, K., Y. Takahashi, T. Berberich, A. Imai, T. Takahashi, A.J. Michael, and T. Kusano. 2007. A protective role for the polyamine spermine against drought stress in *Arabidopsis*. *Biochemical and Biophysical Research Communications* 352:486-490.

Zapata, P.J., M. Serrano, M.T. Pretel, A. Amorós, and M.A. Botella. 2004. Polyamines and ethylene changes during germination of different plant species under salinity. *Plant Science* 167:781-788.

Zeid, I.M., and Z.A. Shedeed. 2006. Response of alfalfa to putrescine treatment under drought stress. *Biologia Plantarum* 50:635-640.


Zhao, F., C. Song, J. He, and H. Zhu. 2007. Polyamines improve K<sup>+</sup>/Na<sup>+</sup> homeostasis in barley seedlings by regulating root ion channel activities. *Plant Physiology* 145:1061-1072.

Zhu, J.K. 2001. Plant salt tolerance. *Trends in Plant Science* 6:66-71.

Zhu, H., G.H. Ding, K. Fang, F.G. Zhao, and P. Qin. 2006. New perspective on the mechanism of alleviating salt stress by spermidine in barley seedlings. *Plant Growth Regulation* 49:147-156.

**Table 1.** *Opuntia* species with seed dormancy.

<b>Species</b>	<b>Authors</b>
<i>O. atrispina</i>	Pendley (2001)
<i>O. auranthiaca</i>	Archibald (1939)
<i>O. camanchica</i>	Pendley (2001)
<i>O. chisoensis</i>	Pendley (2001)
<i>O. compressa</i>	Baskin and Baskin (1977)
<i>O. decumbens</i>	Godínez-Álvarez and Valiente-Banuet (1998)
<i>O. discata</i>	Potter (1984)
<i>O. echios</i> var. <i>gigantea</i>	Wiggins and Focht (1968)
<i>O. edwardsii</i>	Potter et al. (1984)
<i>O. engel</i>	Pendley (2001)
<i>O. engelmannii</i>	Potter et al. (1984); Pendley (2001)
<i>O. ficus-indica</i>	Wang et al. (1996); Altare et al. (2006); D'Aubeterre et al. (2006)
<i>O. humifusa</i>	Cook (1942)
<i>O. imbricata</i>	Pilcher (1970)
<i>O. joconostle</i>	Sánchez-Venegas (1997)
<i>O. kleiniae</i>	Pilcher (1970)
<i>O. leucotricha</i>	Olivares et al. (1999)
<i>O. lindheimeri</i>	Pilcher (1970); Potter et al. (1984)
<i>O. littoralis</i>	Silverstein (2005)
<i>O. macrocentra</i>	Mandujano et al. (2007)
<i>O. macrorhiza</i>	Timmons (1942)
<i>O. microdasys</i>	Mandujano et al. (2007)
<i>O. poly</i>	Pendley (2001)
<i>O. polyacantha</i>	Smreciu et al. (1988)
<i>O. rastrera</i>	Mandujano et al. (1997, 2005)
<i>O. spinossibacca</i>	Pendley (2001)
<i>O. streptacantha</i>	This study
<i>O. stricta</i>	Lotter et al. (1999); Reinhardt et al. (1999)
<i>O. tomentosa</i>	Olvera-Carrillo et al. (2003), Orozco-Segovia et al. (2007)




**Figure 1. 1A)** Germination percentage (Means  $\pm$  SE) of *O. streptacantha* seeds with 1 and 10 mM Pro and with a mixture of PAs (1  $\mu$ M: Put, Spd and Spm).

C = Untreated seeds (Control), PAs (Polyamines) = Put (Putrescine), Spd (Spermidine) and Spm (Spermine) mix, P1 = 1mM Pro (Proline), and P2 = 10 mM Pro (Proline).


**1B)** Germination percentage (Means  $\pm$ SE) with each PAs.

Different letters represent significant differences between treatments.




**Figure 2.** Germination percentage (Means  $\pm$  SE) of *O. streptacantha* seeds under salt stress. C = Untreated seeds (Control), N1 = 25 mM NaCl, N2 = 50 mM NaCl, and N3 = 75 mM NaCl. Different letters represent significant differences between treatments.

A)



B)




**Figure 3.** Germination percentage (Means  $\pm$  SE) of *O. streptacantha* seeds under salt stress with exogenous added 1 mM (Fig. 3A) and 10mM (Fig. 3B) of Pro (Proline).


C = Untreated seeds (Control), P1 = 1 mM Pro, P2 = 10 mM Pro, N1 = 25 mM NaCl, N2 = 50 mM NaCl, and N3 = 75 mM NaCl.

Different letters represent significant differences between treatments.


A)



B)



C)



**Figure 4.** Germination percentage (Means  $\pm$  SE) of *O. streptacantha* seeds under salt stress with exogenous added PAs: Put (Fig. 4A), Spd (Fig. 4B), or Spm (Fig. 4C). C = Untreated seeds (Control), Put = 1  $\mu$ M Putrescine, Spd = 1  $\mu$ M Spermidine, Spm = 1  $\mu$ M Spermine, N1 = 25 mM NaCl, N2 = 50 mM NaCl, and N3 = 75 mM NaCl. Different letters represent significant differences between treatments.